



# Color Glass Condensate and Glasma

BNL, May 2010

## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

## Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

## Matching to hydro

Glasma instabilities  
Hydro in a toy model

## Summary

## Extra bits

François Gelis  
IPhT, CEA/Saclay



## ① Color Glass Condensate

Why small- $x$  gluons matter  
Color Glass Condensate

## ② Just before the collision: Factorization

Stages of AA collisions  
Leading Order  
Leading Log resummation

## ③ Just after the collision: Glasma fields

Initial color fields  
Link to the Lund model  
The ridge in Au-Au collisions

## ④ Matching to hydrodynamics

Glasma instabilities  
Hydro in a toy model

### CGC

Why small- $x$  gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model

### Summary

### Extra bits

**CGC**

Why small-x gluons matter  
Color Glass Condensate

**Factorization**

Stages of AA collisions  
Leading Order  
Leading Logs

**Glasma fields**

Initial color fields  
Link to the Lund model  
Rapidity correlations

**Matching to hydro**

Glasma instabilities  
Hydro in a toy model

**Summary****Extra bits****1 Color Glass Condensate**

Why small-x gluons matter  
Color Glass Condensate

**2 Just before the collision: Factorization**

Stages of AA collisions  
Leading Order  
Leading Log resummation

**3 Just after the collision: Glasma fields**


Initial color fields  
Link to the Lund model  
The ridge in Au-Au collisions

**4 Matching to hydrodynamics**

Glasma instabilities  
Hydro in a toy model

# Longitudinal momentum fraction in AA collisions

## Nucleus-Nucleus collision



- 99% of the multiplicity below  $p_{\perp} \sim 2 \text{ GeV}$
  - $x \sim 10^{-2}$  at RHIC ( $\sqrt{s} = 200 \text{ GeV}$ )
  - $x \sim 4 \cdot 10^{-4}$  at the LHC ( $\sqrt{s} = 5.5 \text{ TeV}$ )
- ▷ partons at small  $x$  are the most important

### CGC

Why small- $x$  gluons matter

Color Glass Condensate

### Factorization

Stages of AA collisions

Leading Order

Leading Logs

### Glasma fields

Initial color fields

Link to the Lund model

Rapidity correlations

### Matching to hydro

Glasma instabilities

Hydro in a toy model

### Summary

### Extra bits



## CGC

Why small-x gluons matter

Color Glass Condensate

## Factorization

Stages of AA collisions

Leading Order

Leading Logs

## Glasma fields

Initial color fields

Link to the Lund model

Rapidity correlations


## Matching to hydro

Glasma instabilities

Hydro in a toy model

## Summary

## Extra bits





## CGC

Why small-x gluons matter

Color Glass Condensate

## Factorization

Stages of AA collisions

Leading Order

Leading Logs

## Glasma fields

Initial color fields

Link to the Lund model

Rapidity correlations

## Matching to hydro

Glasma instabilities

Hydro in a toy model


## Summary

## Extra bits




- Main difficulty: How to treat collisions involving a large number of partons?

# Implications for a QCD approach



- Main difficulty: How to treat collisions involving a large number of partons?
- Dilute regime : one parton in each projectile interact (what the standard perturbative techniques are made for)



- Main difficulty: How to treat collisions involving a large number of partons?
- **Dense regime** : **multiparton processes** become crucial
  - ▷ new techniques are required
  - ▷ multi-parton distributions are needed





CGC = effective theory of small  $x$  gluons

- The **fast partons** ( $k^+ > \Lambda^+$ ) are frozen by time dilation  
▷ described as **static color sources** on the light-cone :

$$J^\mu = \delta^{\mu+} \rho(x^-, \vec{x}_\perp) \quad (0 < x^- < 1/\Lambda^+)$$

- **Slow partons** ( $k^+ < \Lambda^+$ ) cannot be considered static over the time-scales of the collision process  
▷ must be treated as standard gauge fields  
▷ eikonal coupling to the current  $J^\mu$  :  $A_\mu J^\mu$
- The color sources  $\rho$  are **random**, and described by a distribution  $W_{\Lambda^+}[\rho]$ , with  $\Lambda^+$  the longitudinal momentum that separates “soft” and “hard”

## CGC

Why small- $x$  gluons matter

Color Glass Condensate

## Factorization

Stages of AA collisions

Leading Order

Leading Logs

## Glasma fields

Initial color fields

Link to the Lund model

Rapidity correlations

## Matching to hydro

Glasma instabilities

Hydro in a toy model

## Summary

## Extra bits



Independence w.r.t  $\Lambda^+ \rightarrow$  evolution equation (JIMWLK) :

$$\frac{\partial W_{\Lambda^+}}{\partial \ln(\Lambda^+)} = \mathcal{H} W_{\Lambda^+}$$
$$\mathcal{H} = \frac{1}{2} \int_{\vec{x}_\perp, \vec{y}_\perp} \frac{\delta}{\delta \alpha(\vec{y}_\perp)} \eta(\vec{x}_\perp, \vec{y}_\perp) \frac{\delta}{\delta \alpha(\vec{x}_\perp)}$$

where  $-\partial_\perp^2 \alpha(\vec{x}_\perp) = \rho(1/\Lambda^+, \vec{x}_\perp)$

- $\eta(\vec{x}_\perp, \vec{y}_\perp)$  is a non-linear functional of  $\rho$
- Resums all the powers of  $\alpha_s \ln(1/x)$  and of  $Q_s/p_\perp$  that arise in loop corrections
- Simplifies into the BFKL equation when the source  $\rho$  is small (expand  $\eta$  in powers of  $\rho$ )

CGC

Why small-x gluons matter

Color Glass Condensate

Factorization

Stages of AA collisions

Leading Order

Leading Logs

Glasma fields

Initial color fields

Link to the Lund model

Rapidity correlations

Matching to hydro

Glasma instabilities

Hydro in a toy model

Summary

Extra bits



## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

## Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

## Matching to hydro

Glasma instabilities  
Hydro in a toy model

## Summary

## Extra bits

### ① Color Glass Condensate

Why small-x gluons matter  
Color Glass Condensate

### ② Just before the collision: Factorization

Stages of AA collisions  
Leading Order  
Leading Log resummation

### ③ Just after the collision: Glasma fields

Initial color fields  
Link to the Lund model  
The ridge in Au-Au collisions

### ④ Matching to hydrodynamics

Glasma instabilities  
Hydro in a toy model

# Stages of a nucleus-nucleus collision

## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions

Leading Order

Leading Logs

## Glasma fields

Initial color fields

Link to the Lund model

Rapidity correlations


## Matching to hydro

Glasma instabilities

Hydro in a toy model

## Summary

## Extra bits



- The Color Glass Condensate provides a framework to describe nucleus-nucleus collisions up to a time  $\tau \sim Q_s^{-1}$
- Subsequent stages are described as fluid dynamics



## Equations of hydrodynamics :

$$\partial_\mu T^{\mu\nu} = 0$$

## Additional inputs :

EoS :  $p = f(\epsilon)$  , Transport coefficients :  $\eta, \zeta, \dots$

- Required initial conditions :  $T^{\mu\nu}(\tau = \tau_0, \eta, \vec{x}_\perp)$

### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

#### Stages of AA collisions

Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model


### Summary

### Extra bits



# Initial conditions from CGC: power counting

- CGC effective theory with cutoff at the scale  $\Lambda_0^+$  :



$$\mathcal{L} = \underbrace{-\frac{1}{2} \text{tr} F_{\mu\nu} F^{\mu\nu}}_{\text{gluon dynamics}} + \underbrace{(J_1^\mu + J_2^\mu)}_{\text{fast partons}} A_\mu$$

- Expansion in  $g^2$  in the saturated regime:

$$T^{\mu\nu} = \frac{Q_s^4}{g^2} \left[ c_0 + c_1 g^2 + c_2 g^4 + \dots \right]$$

## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions

## Leading Order

Leading Logs

## Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

## Matching to hydro

Glasma instabilities  
Hydro in a toy model

## Summary

## Extra bits

## Initial condition from CGC: Leading Order



### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions

### Leading Order

Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model

### Summary

### Extra bits

- The **Leading Order** contribution is given by **classical fields** :

$$T_{\text{LO}}^{\mu\nu} \equiv c_0 \frac{Q_s^4}{g^2} = \frac{1}{4} g^{\mu\nu} \mathcal{F}^{\lambda\sigma} \mathcal{F}_{\lambda\sigma} - \mathcal{F}^{\mu\lambda} \mathcal{F}^{\nu}_{\lambda}$$


$$\text{with } \underbrace{[\mathcal{D}_\mu, \mathcal{F}^{\mu\nu}]}_{\text{Yang-Mills equation}} = \mathcal{J}^\nu, \quad \lim_{t \rightarrow -\infty} \mathcal{A}^\mu(t, \vec{x}) = 0$$

- The Yang-Mills equations have been solved numerically  
Krasnitz, Venugopalan (1998-2000)  
Lappi (2003)  
Krasnitz, Nara, Venugopalan (2001-2003)



## Initial condition from CGC: Leading Logs

- Consider now quantum corrections to the previous result, restricted to **modes with  $\Lambda_1^+ < k^+ < \Lambda_0^+$**  :



- At leading log accuracy, the contribution of the quantum modes in that strip can be written as :

$$\delta T_{\text{NLO}}^{\mu\nu} = \left[ \ln \left( \frac{\Lambda_0^+}{\Lambda_1^+} \right) \mathcal{H}_1 + \ln \left( \frac{\Lambda_0^-}{\Lambda_1^-} \right) \mathcal{H}_2 \right] T_{\text{LO}}^{\mu\nu}$$

(FG, Lappi, Venugopalan (2008))

### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order

### Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model

### Summary

### Extra bits




## Initial condition from CGC: Leading Logs

- These corrections can be absorbed in the LO result,

$$\left\langle T_{\text{LO}} + \delta T_{\text{NLO}} \right\rangle_{\Lambda_0} = \left\langle T_{\text{LO}} \right\rangle_{\Lambda_1}$$

provided one defines a new effective theory with a lower cutoff  $\Lambda_1^\pm$  and an extended distribution of sources  $W_{\Lambda_1^\pm}[\rho]$ :



$$W_{\Lambda_1^\pm} \equiv \left[ 1 + \ln \left( \frac{\Lambda_0^\pm}{\Lambda_1^\pm} \right) \mathcal{H}_{1,2} \right] W_{\Lambda_0^\pm}$$

(JIMWLK equation for a small change in the cutoff)

### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order

### Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model

### Summary

### Extra bits

## Initial condition from CGC: Leading Logs

- Iterate this step to integrate out all the slow field modes at leading log accuracy:

### Energy-Momentum tensor at Leading Log accuracy

$$\langle T^{\mu\nu}(\tau, \eta, \vec{x}_\perp) \rangle_{\text{LLog}} = \int [D\rho_1 D\rho_2] W_1[\rho_1] W_2[\rho_2] \underbrace{T_{\text{LO}}^{\mu\nu}(\tau, \vec{x}_\perp)}_{\text{for fixed } \rho_{1,2}}$$

- At leading log accuracy, the rapidity dependence comes entirely from the wavefunctions of the projectiles
- This factorization establishes a link to other reactions (such as DIS on a nuclear target) in the saturated regime
- Works for all sufficiently inclusive observables

#### CGC

Why small-x gluons matter  
Color Glass Condensate

#### Factorization

Stages of AA collisions  
Leading Order

#### Leading Logs

#### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

#### Matching to hydro

Glasma instabilities  
Hydro in a toy model

#### Summary

#### Extra bits



## Correlations in $\eta$ and $\vec{x}_\perp$ at Leading Log

- The factorization valid for  $\langle T^{\mu\nu} \rangle$  can be extended to multi-point correlations :

$$\begin{aligned} \langle T^{\mu_1\nu_1}(\tau, \eta_1, \vec{x}_{1\perp}) \cdots T^{\mu_n\nu_n}(\tau, \eta_n, \vec{x}_{n\perp}) \rangle_{\text{LLog}} &= \\ &= \int [D\rho_1 D\rho_2] W_1[\rho_1] W_2[\rho_2] \\ &\quad \times T_{\text{LO}}^{\mu_1\nu_1}(\tau, \vec{x}_{1\perp}) \cdots T_{\text{LO}}^{\mu_n\nu_n}(\tau, \vec{x}_{n\perp}) \end{aligned}$$

- Note: at Leading Log accuracy, all the rapidity correlations come from the evolution of the distributions  $W[\rho_{1,2}]$ 
  - ▷ they are a property of the pre-collision initial state
- This formula predicts long range ( $\Delta\eta \sim \alpha_s^{-1}$ ) rapidity correlations for points located at the same impact parameter

### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order

### Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model

### Summary

### Extra bits



## ① Color Glass Condensate

Why small- $x$  gluons matter  
Color Glass Condensate

## ② Just before the collision: Factorization

Stages of AA collisions  
Leading Order  
Leading Log resummation

## ③ Just after the collision: Glasma fields

Initial color fields  
Link to the Lund model  
The ridge in Au-Au collisions

## ④ Matching to hydrodynamics

Glasma instabilities  
Hydro in a toy model

### CGC

Why small- $x$  gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model


### Summary

### Extra bits

# Initial classical fields, Glasma

Lappi, McLerran (2006)

- Immediately after the collision, the chromo- $\vec{E}$  and  $\vec{B}$  fields are purely longitudinal and boost invariant :



- Glasma** = intermediate stage between the CGC and the quark-gluon plasma

## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

## Glasma fields

### Initial color fields

Link to the Lund model  
Rapidity correlations


## Matching to hydro

Glasma instabilities  
Hydro in a toy model

## Summary

## Extra bits

- The initial chromo- $\vec{E}$  and  $\vec{B}$  fields form longitudinal “flux tubes” extending between the projectiles:



- Correlation length in the transverse plane:  $\Delta r_{\perp} \sim Q_s^{-1}$
- Correlation length in rapidity:  $\Delta \eta \sim \alpha_s^{-1}$
- The flux tubes fill up the entire volume



- A classical field configuration where  $\mathbf{B}_a^i = \lambda \mathbf{E}_a^i$  has an energy-momentum tensor of the form:

$$\langle T^{\mu\nu}(0^+, \eta, \vec{x}) \rangle = \begin{pmatrix} \epsilon & & & \\ & \epsilon & & \\ & & \epsilon & \\ & & & -\epsilon \end{pmatrix}$$

- Multiplicity distribution: Negative Binomial (T. Lappi's talk)
- Long range correlations in rapidity survive in the final state
- $\mathbf{E}$  parallel to  $\mathbf{B} \triangleright$  non-zero  $\tilde{F}F$

## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

## Glasma fields

### Initial color fields

Link to the Lund model  
Rapidity correlations

## Matching to hydro

Glasma instabilities  
Hydro in a toy model

## Summary

## Extra bits

## Link to the Lund string model

- Tanji (2008), Fukushima, FG, Lappi (2009): the yield from the Schwinger mechanism has exactly the same form as the NLO correction in the CGC
- Analogies between the **Glasma** and the **Lund strings**:

Glasma tubes	$\longleftrightarrow$	strings
Negative $P_z$	$\longleftrightarrow$	string tension
Glasma instability	$\longleftrightarrow$	string breaking

- Differences:
  - **$B$**  field in the Glasma
  - The “string size” is set dynamically in the Glasma
  - The Glasma is more closely related to QCD

### CGC

Why small- $x$  gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields

### Link to the Lund model

Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model

### Summary

### Extra bits



# Importance of initial rapidity correlations



## Early physics can survive in long range rapidity correlations

$$t_{\text{correlation}} \leq t_{\text{freeze out}} e^{-\frac{1}{2}|\eta_A - \eta_B|}$$

### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model


### Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model

### Summary

### Extra bits





### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model


### Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model

### Summary

### Extra bits



- Long range correlation in  $\Delta\eta$  (rapidity)
- Narrow correlation in  $\Delta\phi$  (azimuthal angle)


## 2-hadron correlations at RHIC

Dumitru, FG, McLerran, Venugopalan (2008)

Dusling, Fernandez-Fraile, Venugopalan (2009)

Dusling, FG, Lappi, Venugopalan (2009)

- $\eta$ -independent fields lead to long range correlations :



### CGC

Why small-x gluons matter

Color Glass Condensate

### Factorization

Stages of AA collisions

Leading Order

Leading Logs

### Glasma fields

Initial color fields

Link to the Lund model

### Rapidity correlations

### Matching to hydro

Glasma instabilities

Hydro in a toy model


### Summary

### Extra bits

## 2-hadron correlations at RHIC

Dumitru, FG, McLerran, Venugopalan (2008)  
Dusling, Fernandez-Fraile, Venugopalan (2009)  
Dusling, FG, Lappi, Venugopalan (2009)

- $\eta$ -independent fields lead to long range correlations :



- Particles emitted by different flux tubes are not correlated  
▷  $(RQ_s)^{-2}$  sets the strength of the correlation

### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model

### Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model


### Summary

### Extra bits

## 2-hadron correlations at RHIC

Dumitru, FG, McLerran, Venugopalan (2008)  
Dusling, Fernandez-Fraile, Venugopalan (2009)  
Dusling, FG, Lappi, Venugopalan (2009)

- $\eta$ -independent fields lead to long range correlations :



- Particles emitted by different flux tubes are not correlated
  - ▷  $(RQ_s)^{-2}$  sets the strength of the correlation
- At early times, the correlation is flat in  $\Delta\varphi$

### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model

### Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model


### Summary

### Extra bits

## 2-hadron correlations at RHIC

Dumitru, FG, McLerran, Venugopalan (2008)  
Dusling, Fernandez-Fraile, Venugopalan (2009)  
Dusling, FG, Lappi, Venugopalan (2009)

- $\eta$ -independent fields lead to long range correlations :



- Particles emitted by different flux tubes are not correlated
  - ▷  $(RQ_s)^{-2}$  sets the strength of the correlation
- At early times, the correlation is flat in  $\Delta\varphi$   
A collimation in  $\Delta\varphi$  is produced later by radial flow

### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model

### Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model

### Summary

### Extra bits



## ① Color Glass Condensate

Why small- $x$  gluons matter  
Color Glass Condensate

## ② Just before the collision: Factorization

Stages of AA collisions  
Leading Order  
Leading Log resummation

## ③ Just after the collision: Glasma fields

Initial color fields  
Link to the Lund model  
The ridge in Au-Au collisions

## ④ Matching to hydrodynamics

Glasma instabilities  
Hydro in a toy model

### CGC

Why small- $x$  gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model

### Summary

### Extra bits

- Glasma:

$$\langle T^{\mu\nu}(0^+, \eta, \vec{x}) \rangle = \begin{pmatrix} \epsilon & & & \\ & \epsilon & & \\ & & \epsilon & \\ & & & -\epsilon \end{pmatrix}$$

- Ideal hydro:

$$T_{\text{ideal}}^{\mu\nu}(0^+, \eta, \vec{x}) = \begin{pmatrix} \epsilon & & & \\ & p & & \\ & & p & \\ & & & p \end{pmatrix}$$

- If a smooth matching from the Glasma to Hydro is possible, one should be able to recover the fluid behavior from classical fields



## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

## Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

## Matching to hydro

Glasma instabilities  
Hydro in a toy model


## Summary

## Extra bits



Romatschke, Venugopalan (2005)

- Perturbations to the classical fields grow like  $\exp(\sqrt{Q_s \tau})$  until the non-linearities become important :



- ▷ Quantum fluctuations become  $\mathcal{O}(1)$  corrections when

$$\tau \sim \tau_{\max} \sim Q_s^{-1} \ln^2(1/\alpha_s)$$



## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

## Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

## Matching to hydro

Glasma instabilities  
Hydro in a toy model

## Summary

## Extra bits

## Resummation of the unstable terms

- To go beyond the time  $\tau_{\max}$ , one must resum all the **fastest growing terms**  $\sim [g^2 e^{\sqrt{Q_s \tau}}]^n$
- This amounts to superimposing fluctuations to the initial classical field:

$$\begin{aligned} \langle T^{\mu\nu}(\tau, \eta, \vec{x}_\perp) \rangle &\stackrel{\text{LLog resummed}}{=} \int [D\rho_1 D\rho_2] \, W_{Y_1}[\rho_1] W_{Y_2}[\rho_2] \\ &\times \int [Da] \, F[a] \, T_{\text{LO}}^{\mu\nu}[\underbrace{\mathcal{A} + a}_{\text{initial field}}] \end{aligned}$$

- Fukushima, FG, McLerran (2006)**: this result can be obtained in a semi-classical approach (with Gaussian  $F[a]$ )
- FG, Lappi, Venugopalan (2008)**: can be obtained by a resummation of the NLO result in the CGC

### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

### Matching to hydro

Glasma instabilities  
Hydro in a toy model

### Summary

### Extra bits



## Hydro behavior in a toy model

- Including the fluctuations in the calculation of  $T^{\mu\nu}$  is hard:
  - Space-dependent fluctuations need to be renormalized (because of UV divergences)
  - The QCD classical equations are difficult to solve

### Toy model: scalar field, uniform in $\eta$

Equation of motion: 
$$\ddot{\phi} + \frac{1}{\tau}\dot{\phi} - \nabla_{\perp}^2\phi + V'(\phi) = 0$$

Interaction potential: 
$$V(\phi) \sim g^2\phi^4$$

Initial conditions at  $\tau = \tau_0$ : 
$$\phi = \varphi_0, \quad \dot{\phi} = \dot{\varphi}_0$$

Gaussian fluctuations of  $\varphi_0$  and  $\dot{\varphi}_0$

#### CGC

Why small-x gluons matter  
Color Glass Condensate

#### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

#### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

#### Matching to hydro

Glasma instabilities

#### Hydro in a toy model


#### Summary

#### Extra bits



# Hydro behavior in a toy model (1+0 dim)

## Classical field evolution w/o fluctuations



### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

### Matching to hydro

Glasma instabilities

### Hydro in a toy model

### Summary


### Extra bits

- Without fluctuations,  $p$  oscillates forever



# Hydro behavior in a toy model (1+0 dim)

## Classical field evolution with fluctuations



### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

### Matching to hydro

Glasma instabilities

### Hydro in a toy model

### Summary


### Extra bits

- Without fluctuations,  $p$  oscillates forever
- With fluctuations,  $p$  relaxes quickly to  $\epsilon/3$



# Hydro behavior in a toy model (1+0 dim)

## Classical field evolution with fluctuations



### CGC

Why small- $x$  gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

### Matching to hydro

Glasma instabilities

### Hydro in a toy model

### Summary

### Extra bits

- Without fluctuations,  $p$  oscillates forever
- With fluctuations,  $p$  relaxes quickly to  $\epsilon/3$
- $\epsilon$  and  $p$  decrease as  $1/\tau^{4/3}$ 
  - ▷ same behavior as in ideal hydro with EoS  $\epsilon = 3p...$



## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

## Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

## Matching to hydro

Glasma instabilities

## Hydro in a toy model


## Summary

## Extra bits

# Hydro behavior in a toy model (1+2 dim)

- Left:  $\log |(\epsilon - 3p)/\epsilon|$  without fluctuations
- Right:  $\log |(\epsilon - 3p)/\epsilon|$  with fluctuations

time = 0.1





## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

## Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

## Matching to hydro

Glasma instabilities

## Hydro in a toy model


## Summary

## Extra bits

# Hydro behavior in a toy model (1+2 dim)

- Left:  $\log |(\epsilon - 3p)/\epsilon|$  without fluctuations
- Right:  $\log |(\epsilon - 3p)/\epsilon|$  with fluctuations

time = 0.35







## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

## Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

## Matching to hydro

Glasma instabilities

## Hydro in a toy model


## Summary

## Extra bits

# Hydro behavior in a toy model (1+2 dim)

- Left:  $\log |(\epsilon - 3p)/\epsilon|$  without fluctuations
- Right:  $\log |(\epsilon - 3p)/\epsilon|$  with fluctuations

time = 1.20





## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

## Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

## Matching to hydro

Glasma instabilities

## Hydro in a toy model


## Summary

## Extra bits

# Hydro behavior in a toy model (1+2 dim)

- Left:  $\log |(\epsilon - 3p)/\epsilon|$  without fluctuations
- Right:  $\log |(\epsilon - 3p)/\epsilon|$  with fluctuations

time = 4.16





## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

## Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

## Matching to hydro

Glasma instabilities

## Hydro in a toy model


## Summary

## Extra bits

# Hydro behavior in a toy model (1+2 dim)

- Left:  $\log |(\epsilon - 3p)/\epsilon|$  without fluctuations
- Right:  $\log |(\epsilon - 3p)/\epsilon|$  with fluctuations



time = 14.43



# Hydro behavior in a toy model (1+2 dim)

- Left: magnitude of the viscous tensor  $\log(\Pi^{\mu\nu}/\epsilon)$
- Right: velocity field

time = 14.43



## Initial state, up to $\tau = 0^+$

- Consistent framework to include the non-linear saturation effects in heavy ion collisions
- Factorization of the large logs of  $x_{1,2}$  into universal distributions  $W[\rho]$
- Implies long range rapidity correlations, in good agreement with RHIC data

## Final state evolution

- Unstable fluctuations need to be resummed, but the machinery for doing that is not fully developed
- An equation of state may be obtained by superimposing quantum fluctuations to the classical fields, without complete thermalization of the system

### CGC

Why small-x gluons matter  
Color Glass Condensate

### Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

### Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

### Matching to hydro


Glasma instabilities  
Hydro in a toy model

### Summary

### Extra bits



$$\mathcal{L} \equiv \frac{1}{2} [\dot{\phi}_1^2 + \dot{\phi}_2^2] - \frac{g^2}{4!} [\phi_1^2 + \phi_2^2]^2$$



## CGC

Why small-x gluons matter  
Color Glass Condensate

## Factorization

Stages of AA collisions  
Leading Order  
Leading Logs

## Glasma fields

Initial color fields  
Link to the Lund model  
Rapidity correlations

## Matching to hydro

Glasma instabilities  
Hydro in a toy model

## Summary

## Extra bits